

Higgs mechanism

The *Higgs mechanism* enables the massless gauge bosons in a gauge theory to acquire effective mass through interaction with auxiliary fields (called the Higgs fields). The mechanism is based on the phenomenon of spontaneous symmetry breaking.

In the standard model the Higgs mechanism generates the masses of the W^\pm and Z weak gauge bosons.

Spontaneous symmetry breaking

Spontaneous symmetry breaking takes place when the Lagrangian of a system is symmetric while the ground state isn't.

Here is the canonical example: a complex self-interacting scalar (Higgs) field φ with a globally $U(1)$ symmetric Lagrangian,

$$\mathcal{L} = \partial_a \varphi^* \partial^a \varphi - V_{\text{MH}}(\varphi^* \varphi) \quad (1)$$

where the potential energy V_{MH} has the ‘‘Mexican hat’’ form (see fig. 1),

$$V_{\text{MH}}(\varphi^* \varphi) = \frac{\lambda^2}{4} (\varphi^* \varphi - m^2)^2, \quad (2)$$

where λ is a dimensionless constant and m is a constant with the dimension of mass¹.




Figure 1: ‘‘Mexican hat’’ potential (2) in arbitrary units as function of the real, $\Re\varphi$, and imaginary, $\Im\varphi$, parts of the complex scalar field φ .

The lowest energy state of the system is reached for a finite (non-zero) field φ_0 (often called Higgs condensate or vacuum expectation value) such that $\varphi_0^* \varphi_0 = m^2$. Due to the symmetry of the Lagrangian there are infinitely many vacuum states $\varphi_0 = m e^{i\xi}$, where $0 \leq \xi < 2\pi$.

¹hypothetical particles with negative square mass are called *tachyons*.

Breaking the symmetry means choosing a particular vacuum state, e.g. $\varphi_0 = m$. Let us look at small fluctuations around this vacuum state in the form

$$\varphi = (m + \eta)e^{i\theta/m}. \quad (3)$$

In terms of the new fields η and θ the Lagrangian becomes (up to the second order terms)

$$\mathcal{L} = \partial_a \eta \partial^a \eta - \lambda^2 m^2 \eta^2 + \partial_a \theta \partial^a \theta. \quad (4)$$

This Lagrangian apparently describes a massless boson θ (called *Goldstone boson*) and a massive boson η (to be called Higgs boson in the standard model) with the (positive) mass $m_H = \lambda m$.

This simple model actually illustrates the *Goldstone theorem* which states that whenever a continuous symmetry is spontaneously broken, a massless (or light, if the symmetry was not exact) scalar particle appears in the spectrum of the system. It corresponds to excitations along the ‘‘symmetry direction’’.

Higgs mechanism

Higgs mechanism is spontaneous symmetry breaking in a gauge theory.

Let us make our $U(1)$ symmetric model (1) gauge invariant by coupling our Higgs field φ to a real gauge field B_a ,

$$\mathcal{L} = D_a^* \varphi^* D^a \varphi - V_{\text{MH}}(\varphi^* \varphi) + \mathcal{L}_{\text{YM}}, \quad (5)$$

where $D_a = \partial_a + igB_a$ is the gauge-group covariant derivative, and

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{ab} F^{ab}, \quad F_{ab} = \frac{1}{ig} [D_a, D_b] \quad (6)$$

is the Yang-Mills Lagrangian of the free gauge field B_a .

Again let us look for small fluctuations of the Higgs field from the condensate, $\varphi = (m + \eta)e^{i\theta/m}$. The covariant derivative,

$$D_a \phi = (\partial_a + igB_a)(m + \eta)e^{i\theta/m}, \quad (7)$$

with up to linear terms in the fields is given as

$$D_a \phi \approx (\partial_a \eta + i\partial_a \theta + igmB_a)e^{i\theta/m}. \quad (8)$$

Introducing the new field

$$B'_a = B_a + \frac{1}{gm} \partial_a \theta, \quad (9)$$

the derivative becomes

$$D_a \phi \approx (\partial_a \eta + igmB'_a)e^{i\theta/m}. \quad (10)$$

Now the kinetic term in the Lagrangian (5) becomes where

$$D_a^* \varphi^* D_a \varphi \approx \partial_a \eta \partial^a \eta + g^2 m^2 B'^a B'_a. \quad (11)$$

It is easy to see that this is actually a result of the gauge transformation

$$\begin{cases} \varphi & \rightarrow \varphi' = e^{-i\theta/m} \varphi \\ B_a & \rightarrow B'_a = B_a + \frac{1}{gm} \partial_a \theta. \end{cases} \quad (12)$$

of the original fields $\varphi = (m + \eta)e^{i\theta/m}$ and B_a .

Indeed,

$$D'_a \varphi' = (\partial_a + ig B'_a)(m + \eta) \approx \partial_a \eta + igm B'_a, \quad (13)$$

and

$$D_a^* \varphi'^* D'_a \varphi' \approx \partial_a \eta \partial^a \eta + g^2 m^2 B'^a B'_a. \quad (14)$$

Finally the total Lagrangian becomes

$$\mathcal{L} = \partial_a \eta \partial^a \eta - \lambda^2 m^2 \eta^2 + \mathcal{L}_{\text{YM}} + g^2 m^2 B'^a B'_a + (\text{interaction terms}). \quad (15)$$

This Lagrangian describes a massive scalar Higgs boson η and a massive vector boson B_a .

The mass of the boson, $m_B = gm = g\varphi_0$, is determined by the Higgs condensate ϕ_0 and the coupling g between the Higgs and the gauge fields.

Before the symmetry breaking there were 4 degrees of freedom: two in the complex Higgs field φ and two in the real massless (and hence transverse) gauge field B_a .

After the symmetry breaking there appeared a massive real scalar boson η (Higgs boson); the gauge boson acquired a finite mass gm and therefore lost its transverse character; the θ field has turned into the longitudinal polarization component of the gauge field and disappeared from the Lagrangian: in total again 4 degrees of freedom.

The standard model

The standard model is a Yang-Mills field theory with the symmetry $U(1) \times SU(2) \times SU(3)$. It consists of the electroweak sector², $U(1) \times SU(2)_L$; the strong sector, $SU(3)$, called quantum chromodynamics (QCD); and the Higgs sector.

The QCD sector

The QCD Lagrangian is an ordinary Yang-Mills Lagrangian with $SU(3)$ symmetry,

$$\mathcal{L}_{\text{QCD}} = \sum_q \bar{q} (i\gamma^a D_a - m_q) q + \mathcal{L}_{\text{YM}} \quad (16)$$

²the subscript L means that only left bispinors take part in the interaction.

$$D_a = \partial_a + ig_s I^j A_a^j, \quad (17)$$

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{ab}^j F_j^{ab}, \quad (18)$$

$$F_{ab}^j = \partial_a A_b - \partial_b A_a - g_s f_{kl}^j A_a^k A_b^l, \quad (19)$$

where $q \in \{u, d, s, c, t, b\}$ is the quark field of specific flavour (Lorentz bispinor and color triplet); I^j are the generators of the $SU(3)$ group, usually represented as $\frac{1}{2}\lambda^j$ where λ^j are Gell-Mann matrices; f_{jkl} are the structure constants of the $SU(3)$ group; and g_s is the QCD coupling constant.

Electroweak sector

The electroweak sector is a "weakly mixed" $U(1) \times SU(2)_L$,

$$\begin{aligned} \mathcal{L}_{\text{EW}} = \sum_{\psi} \bar{\psi} \gamma^a \left(i\partial_a - g' \frac{Y_W}{2} B_a - g \vec{T}_L \vec{W}_a \right) \psi \\ + \mathcal{L}_{\text{YM}}(B_a) + \mathcal{L}_{\text{YM}}(\vec{W}_a), \end{aligned} \quad (20)$$

where B_a is the $U(1)$ gauge field; \vec{W}_a is the three-component $SU(2)$ gauge field; g' and g are coupling constants; Y_W is the *weak hyper-charge* – the generator of the $U(1)$ group; $\vec{T} = \frac{1}{2}\vec{\tau}$ (where $\vec{\tau}$ are the Pauli matrices in the weak isospin $SU(2)$ space) are the infinitesimal generators of the $SU(2)$ group, subscript L indicates that they only act on the left fermions ψ_L , which are weak doublets while the right fermions ψ_R are weak singlets. For example, for the leptons of the first generation,

$$\psi_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}, \quad \psi_R = e_R. \quad (21)$$

The second singlet, the right neutrino, if exists, is "sterile" (if massless) as it does not interact at all. For the quarks of the first generation,

$$\psi_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix}, \quad \psi_R = u_R, d_R. \quad (22)$$

The weak hypercharge Y_W is defined such that the electric charge Q is equal

$$Q = (T_L)_3 + \frac{1}{2} Y_W. \quad (23)$$

The gauge field that interacts with the third component of the weak isospin τ_3 but does not interact with the electric charge Q is called Z -boson,

$$Z_a = \frac{gW_a^{(3)} - g'B_a}{\sqrt{g^2 + g'^2}}. \quad (24)$$

The orthogonal combination, which interacts with Q but not with τ_3 , becomes the electromagnetic field A_a ,

$$A_a = \frac{g'W_a^{(3)} + gB_a}{\sqrt{g^2 + g'^2}}. \quad (25)$$

Introducing the *weak mixing angle*³, $\tan \theta_W = g'/g$, the transformation can be conveniently written as an orthogonal transformation,

$$\begin{pmatrix} A \\ Z \end{pmatrix} = \begin{pmatrix} \cos \theta_W & \sin \theta_W \\ -\sin \theta_W & \cos \theta_W \end{pmatrix} \begin{pmatrix} B \\ W^3 \end{pmatrix}, \quad (26)$$

which apparently preserves the "kinetic energy" terms in the Lagrangian.

With the notations $e = g \sin \theta_W$, $W^\pm = \frac{1}{\sqrt{2}}(W^1 \mp iW^2)$, $T^\pm = \frac{1}{2}(\tau^1 \pm i\tau^2)$ the fermionic part of the Lagrangian takes the standard form,

$$\begin{aligned} \mathcal{L}_{\text{EW}}^{(\text{F})} &= \sum_{\psi} \bar{\psi} \gamma^a i \partial_a \psi - e \sum_{\psi} \bar{\psi} \gamma^a Q \psi A_a \\ &- \frac{g}{\cos \theta_W} \sum_{\psi} \bar{\psi} \gamma^a \left(\frac{1}{2} \tau_L^3 - Q \sin^2 \theta_W \right) \psi Z_a \\ &- \frac{g}{2\sqrt{2}} \sum_{\psi} \bar{\psi} \gamma^a (T_L^+ W^+ + T_L^- W^-) \psi. \end{aligned} \quad (27)$$

Higgs sector

The Higgs field must provide masses to three gauge fields – hence, together with the Higgs boson, there should be minimum 4 components in the Higgs field. In the standard model it is a complex spinor of the group $SU(2)_L$,

$$\varphi = \frac{1}{\sqrt{2}} \begin{pmatrix} \varphi^+ \\ \varphi^0 \end{pmatrix}, \quad (28)$$

where the indexes $+$ and 0 indicate the Q -charges of the components; the Y_W -charge of both components is equal 1.

The Higgs sector Lagrangian is

$$\begin{aligned} \mathcal{L}_H &= \varphi^\dagger \left(\overleftarrow{\partial}_a - ig' \frac{1}{2} Y_W B_a - ig \frac{1}{2} \vec{\tau} \vec{W}_a \right) \cdot \\ &\quad \cdot \left(\overrightarrow{\partial}_a + ig' \frac{1}{2} Y_W B_a + ig \frac{1}{2} \vec{\tau} \vec{W}_a \right) \varphi \\ &\quad - \frac{\lambda^2}{4} (\varphi^\dagger \varphi - v^2)^2 \end{aligned} \quad (29)$$

After the spontaneous symmetry breaking the fields Z and W^\pm get masses, $m_w^2 = \frac{1}{4}v^2g^2$, $m_Z^2 \cos^2 \theta_w = m_w^2$, while the field A remains massless.

³also called Weinberg angle.