

### Finite rotation matrix

The generators can recover not only infinitesimal,  $g(d\alpha) = 1 + iI d\alpha$ , but (in principle) also finite elements of the group (with certain caveats). It is especially easy with *additive parameters*, where  $g(\alpha + d\alpha) = g(\alpha)g(d\alpha)$ . Indeed, in this case

$$g(\alpha + d\alpha) = g(d\alpha)g(\alpha) = (1 + iI d\alpha)g(\alpha), \quad (1)$$

and thus the group element  $g(\alpha)$  satisfies the differential equation

$$\frac{\partial g(\alpha)}{\partial \alpha} = iI g(\alpha), \quad (2)$$

with the initial condition  $g(0) = 1$ . One can check by direct substitution that the solution to this equation is exponential function, understood as an infinite Taylor series

$$g(\alpha) = e^{iI\alpha} \equiv \sum_{n=0}^{\infty} \frac{(iI\alpha)^n}{n!}. \quad (3)$$

For example, for the rotation around a given axis  $\vec{n}$  the rotation angle  $\theta$  is an additive parameter of the rotation matrix  $R(\vec{n}, \theta)$ ,

$$\begin{aligned} R(\vec{n}, \theta + d\theta) &= R(\vec{n}, \theta)R(\vec{n}, d\theta) \\ &= (1 + i\vec{I} \cdot \vec{n} d\theta)R(\vec{n}, \theta), \end{aligned} \quad (4)$$

and thus the rotation matrix satisfies the differential equation

$$\frac{dR}{d\theta} = i\vec{I} \cdot \vec{n} R, \quad (5)$$

with the boundary condition  $R(\vec{n}, 0) = 1$ . The solution is given by the Taylor series,

$$R(\vec{n}, \theta) = e^{i\vec{I} \cdot \vec{n} \theta} \equiv 1 + i\vec{I} \cdot \vec{n} \theta + \frac{(i\vec{I} \cdot \vec{n} \theta)^2}{2!} + \dots \quad (6)$$

In practice for a finite-dimension representation there is only a finite number of terms in the series.

### Direct product of two representations

In the field theory we often have to work with different types of products of covariant quantities, like  $\partial_a j^a$  or  $\partial_a A^b$ . We need to know how such products transform.

For simplicity we shall only consider direct products and direct sums of matrices, although the concepts are also defined for abstract groups.

A *direct product*  $g \otimes h$  of two matrices  $g$  and  $h$  is a matrix made of all possible (and suitable arranged) pairwise products of matrix elements of  $g$  and  $h$ ,

$$g \otimes h = \begin{bmatrix} g_{11}h_{11} & \dots & \dots \\ \dots & \dots & \dots \\ \dots & \dots & g_{n_g n_g} h_{n_h n_h} \end{bmatrix}, \quad (7)$$

where  $n_g$  and  $n_h$  are the sizes of  $g$  and  $h$ . The size of the direct product is  $n_{g \otimes h} = n_g n_h$ .

If the matrices  $g$  and  $h$  act on column-vectors  $v_g$  and  $v_h$  of sizes  $n_g$  and  $n_h$ , then the matrices  $g \otimes h$  act on a column-vector  $v_g \otimes v_h$  of size  $n_g n_h$  consisting of all pairwise products of the elements of vectors  $v_g$  and  $v_h$ ,

$$v_g \otimes v_h = \begin{bmatrix} (v_g)_1 (v_h)_1 \\ \dots \\ (v_g)_{n_g} (v_h)_{n_h} \end{bmatrix}. \quad (8)$$

A *direct sum*  $g \oplus h$  of two matrices  $g$  and  $h$  is a block-diagonal matrix made of matrices  $g$  and  $h$ ,

$$g \oplus h = \begin{bmatrix} g & 0 \\ 0 & h \end{bmatrix}. \quad (9)$$

The size of the direct sum is  $n_{g \oplus h} = n_g + n_h$ .

If the matrices  $g$  and  $h$  act on column-vectors  $v_g$  and  $v_h$  of sizes  $n_g$  and  $n_h$ , then the matrices  $g \oplus h$  act on a column-vector  $v_g \oplus v_h$  of size  $n_g + n_h$  which is made of elements of both vectors  $v_g$  and  $v_h$ ,

$$v_g \oplus v_h = \begin{bmatrix} v_g \\ v_h \end{bmatrix}. \quad (10)$$

Suppose we have two (different) representations,  $G = \{g\}$  and  $H = \{h\}$ , of some Lie group  $\{\Lambda\}$ , with the corresponding infinitesimal elements<sup>1</sup>

$$g = 1_G + i\vec{I}_G \vec{\alpha}, \quad (11)$$

$$h = 1_H + i\vec{I}_H \vec{\alpha}, \quad (12)$$

where  $\vec{\alpha}$  are the parameters of the group  $\{\Lambda\}$ .

It is easy to show that the generators  $I_{G \otimes H}$  of a direct product  $G \otimes H$  of two representations  $G$  and  $H$  of the same group is a (Kronecker) sum of the corresponding generators  $I_G$  and  $I_H$ ,

$$\vec{I}_{G \otimes H} = \vec{I}_G \otimes 1_H + 1_G \otimes \vec{I}_H. \quad (13)$$

Indeed,

$$\begin{aligned} g \otimes h &= (1_G + i\vec{I}_G \vec{\alpha}) \otimes (1_H + i\vec{I}_H \vec{\alpha}) \\ &= 1_G \otimes 1_H + i(\vec{I}_G \otimes 1_H + 1_G \otimes \vec{I}_H) \vec{\alpha} \\ &= 1 + i\vec{I}_{G \otimes H} \vec{\alpha}, \end{aligned} \quad (14)$$

$$\Rightarrow \vec{I}_{G \otimes H} = \vec{I}_G \otimes 1_H + 1_G \otimes \vec{I}_H. \quad (15)$$

### Clebsch-Gordan theorem

The direct product  $(j_1) \otimes (j_2)$  of two irreducible representations of the rotation group is a reducible

<sup>1</sup> $\vec{I} \vec{\alpha} \equiv \sum_k I_k \alpha_k$

representation which can be reduced into a direct sum of irreducible representations,

$$(j_1) \otimes (j_2) = \sum_{j=|j_1-j_2|}^{j_1+j_2} \oplus(j). \quad (16)$$

*Example:* a direct product  $\vec{a} \otimes \vec{b}$  of two vectors,  $(1) \otimes (1)$ , reduces to a direct sum of a scalar ( $j = 0$ ), an antisymmetric tensor ( $j = 1$ ), and a symmetric tensor with zero trace ( $j = 2$ ),

$$\vec{a} \otimes \vec{b} = (\vec{a}\vec{b}) \oplus (\vec{a} \times \vec{b}) \oplus \left( a_i b_j + a_j b_i - \frac{2}{3}(\vec{a}\vec{b})\delta_{ij} \right). \quad (17)$$

### Irreducible representations of the Lorentz group

With the complex parameterization

$$d\vec{w} = \vec{n}d\theta + id\vec{v} \quad (18)$$

the infinitesimal Lorentz transformation is given as

$$\Lambda = 1 + i\vec{M}d\vec{w} + i\vec{N}d\vec{w}^*, \quad (19)$$

where the generators  $M$  and  $N$  satisfy the Lie algebra

$$M_k M_l - M_l M_k = i \sum_m \epsilon_{klm} M_m \quad (20)$$

$$N_k N_l - N_l N_k = i \sum_m \epsilon_{klm} N_m \quad (21)$$

$$M_k N_l - N_l M_k = 0, \quad (22)$$

that is, two independent rotation Lie algebras. An infinitesimal matrix  $t$  from a representation of the Lorentz group is then given as

$$t = \mathbf{1}_M \otimes \mathbf{1}_N + i\vec{M} \otimes \mathbf{1}_N d\vec{w} + i\mathbf{1}_M \otimes \vec{N} d\vec{w}^*, \quad (23)$$

where  $\mathbf{1}_M$  and  $\mathbf{1}_N$  are the unit matrices in the spaces of  $M$  and  $N$  generators.

Thus an irreducible representation of the Lorentz group is determined by two numbers  $(m, n)$ , each taking non-negative integer or half-integer values. The dimensions of the  $M$  and  $N$ -generators are then  $(2m + 1)$  and  $(2n + 1)$  correspondingly. The dimension of the representation is  $(2m + 1)(2n + 1)$ .

**Direct product of two irreducible representations** There exists a similar theorem for the Lorentz group,

$$(j_1, k_1) \otimes (j_2, k_2) = \sum_{j=|j_1-j_2|}^{j_1+j_2} \sum_{k=|k_1-k_2|}^{k_1+k_2} \oplus(j, k). \quad (24)$$

**Rotational properties of an irreducible representation of the Lorentz group** If we only consider rotations,  $d\vec{v} = 0$ , the infinitesimal element of a Lorentz group representation (23) becomes

$$g|_{d\vec{v}=0} = 1 + i \left( \vec{M} \otimes \mathbf{1}_N + \mathbf{1}_M \otimes \vec{N} \right) \vec{n}d\theta, \quad (25)$$

which can be identified as the Kronecker sum (20) of two generators with rotation Lie algebra. A Kronecker sum of generators corresponds to a direct product of their representations.

Thus, under rotations, an irreducible representations  $(j_1, j_2)$  of the Lorentz group reduces to a direct sum of irreducible representations of the rotation group ( $j$ ) with  $j = |j_1 - j_2|, \dots, j_1 + j_2$ .

*Example:* a four-vector  $\{E, \mathbf{p}\}$  transforms under  $(\frac{1}{2}, \frac{1}{2})$  representation of the Lorentz group and under rotations reduces to a direct sum of a scalar  $E$  and a vector  $\mathbf{p}$ .

**Parity transformation and irreducible representations of the Lorentz group** Parity transformation is the simultaneous change of spatial coordinates,

$$P \begin{pmatrix} t \\ \vec{x} \end{pmatrix} = \begin{pmatrix} t \\ -\vec{x} \end{pmatrix}. \quad (26)$$

The parity transformation, like rotations, reflects our freedom in choosing frames of reference for a description of physical systems and therefore must be included in the group of coordinate transformations in the principle of covariance.

Under the parity transformation the rotation generators do not change,  $\vec{J} \rightarrow \vec{J}$ , while the velocity-boost-generators change sign,  $\vec{K} \rightarrow -\vec{K}$ . The "optimal" generators  $\vec{M} = \frac{1}{2}(\vec{J} - i\vec{K})$  and  $\vec{N} = \frac{1}{2}(\vec{J} + i\vec{K})$  transform into each other,  $\vec{M} \leftrightarrow \vec{N}$ .

Thus an irreducible representation of the Lorentz group  $(j_1, j_2)$  transforms under parity transformation into a representation  $(j_2, j_1)$ ,

$$(j_1, j_2) \xleftrightarrow{\mathcal{P}} (j_2, j_1). \quad (27)$$

Consequently, if  $j_1 \neq j_2$ , the representation of the covariance group of coordinate transformations has to be enlarged to the direct sum  $(j_1, j_2) \oplus (j_2, j_1)$ .